



Enumerating the edge-colourings and total colourings of a regular graph

Stéphane Bessy, Frédéric Havet

► To cite this version:

Stéphane Bessy, Frédéric Havet. Enumerating the edge-colourings and total colourings of a regular graph. 2012 Workshop on Graph Theory and Combinatorics, Aug 2012, National Sun Yat-sen University, Kaohsiung, Taiwan. lirmm-00811571

HAL Id: lirmm-00811571

<https://hal-lirmm.ccsd.cnrs.fr/lirmm-00811571>

Submitted on 10 Apr 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Enumerating the edge-colourings of a regular graph

Stéphane Bessy and Frédéric Havet

Assistant Professor
LIRMM, UM2, Montpellier

Researcher
CNRS, Sophia-Antipolis

FRANCE

2012 Workshop on Graph Theory and Combinatorics
Department of Applied Mathematics, National Sun Yat-sen University,
Kaohsiung, Taiwan.

Outline

- 1 Introduction
 - Colorings
 - Algorithmic problems
 - Our results
- 2 Enumerating the 3-edge colorings of a cubic graph
 - The 3-edge colorings of a 3-regular graph
 - Turning the proof into algorithm
- 3 Extensions: k -edge colorings and the total colorings
 - k -edge colorings
 - A more precise bounds for the 3-edge coloring
 - Total coloring
- 4 Conclusion

Introduction

- 1 Introduction
 - Colorings
 - Algorithmic problems
 - Our results
- 2 Enumerating the 3-edge colorings of a cubic graph
 - The 3-edge colorings of a 3-regular graph
 - Turning the proof into algorithm
- 3 Extensions: k -edge colorings and the total colorings
 - k -edge colorings
 - A more precise bounds for the 3-edge coloring
 - Total coloring
- 4 Conclusion

Vertex Coloring


Definition (k -vertex coloring)

A k -vertex coloring of a (non oriented) graph $G = (V, E)$ is a function $c : V \rightarrow \{1, \dots, k\}$ such that $uv \in E$ implies $c(u) \neq c(v)$.

Vertex Coloring

Definition (k -vertex coloring)


A k -vertex coloring of a (non oriented) graph $G = (V, E)$ is a function $c : V \rightarrow \{1, \dots, k\}$ such that $uv \in E$ implies $c(u) \neq c(v)$.



Vertex Coloring

Definition (k -vertex coloring)


A k -vertex coloring of a (non oriented) graph $G = (V, E)$ is a function $c : V \rightarrow \{1, \dots, k\}$ such that $uv \in E$ implies $c(u) \neq c(v)$.



Vertex Coloring

Definition (k -vertex coloring)

A k -vertex coloring of a (non oriented) graph $G = (V, E)$ is a function $c : V \rightarrow \{1, \dots, k\}$ such that $uv \in E$ implies $c(u) \neq c(v)$.



Definition (chromatic number)

$\chi(G)$ is the minimum k such that G admits a k -vertex coloring.

Edge Coloring


Definition (k -edge coloring)

A k -edge coloring of a graph $G = (V, E)$ is a function $c : E \rightarrow \{1, \dots, k\}$ such that $uv \in E$ and $uw \in E$, with $v \neq w$ implies $c(uv) \neq c(uw)$.

Edge Coloring

Definition (k -edge coloring)


A k -edge coloring of a graph $G = (V, E)$ is a function $c : E \rightarrow \{1, \dots, k\}$ such that $uv \in E$ and $uw \in E$, with $v \neq w$ implies $c(uv) \neq c(uw)$.



Edge Coloring

Definition (k -edge coloring)


A k -edge coloring of a graph $G = (V, E)$ is a function $c : E \rightarrow \{1, \dots, k\}$ such that $uv \in E$ and $uw \in E$, with $v \neq w$ implies $c(uv) \neq c(uw)$.



Edge Coloring

Definition (k -edge coloring)

A k -edge coloring of a graph $G = (V, E)$ is a function $c : E \rightarrow \{1, \dots, k\}$ such that $uv \in E$ and $uw \in E$, with $v \neq w$ implies $c(uv) \neq c(uw)$.





Definition (chromatic index)

$\chi'(G)$ is the minimum k such that G admits a k -edge coloring.



Edge Coloring

- The *line graph* $L(G)$ of G :
 - The vertex set of $L(G)$ is the edge set of G
 - Two edges e and f of G are adjacent in $L(G)$ if there are incident in G

 G  $L(G)$

Edge Coloring

- The *line graph* $L(G)$ of G :
 - The vertex set of $L(G)$ is the edge set of G
 - Two edges e and f of G are adjacent in $L(G)$ if there are incident in G

 G  $L(G)$

- $\chi'(G) = \chi(L(G))$

Total Coloring

Definition (k -total coloring)

A k -total coloring of $G = (V, E)$ is $c : V \cup E \rightarrow \{1, \dots, k\}$ such that:


- $uv \in E$ implies $c(u) \neq c(v)$
- $uv \in E$ and $uw \in E$, with $v \neq w$ implies $c(uv) \neq c(uw)$
- $uv \in E$ implies $c(uv) \neq c(u)$

Total Coloring

Definition (k -total coloring)

A k -total coloring of $G = (V, E)$ is $c : V \cup E \rightarrow \{1, \dots, k\}$ such that:

- $uv \in E$ implies $c(u) \neq c(v)$
- $uv \in E$ and $uw \in E$, with $v \neq w$ implies $c(uv) \neq c(uw)$
- $uv \in E$ implies $c(uv) \neq c(u)$




Total Coloring

Definition (k -total coloring)

A k -total coloring of $G = (V, E)$ is $c : V \cup E \rightarrow \{1, \dots, k\}$ such that:

- $uv \in E$ implies $c(u) \neq c(v)$
- $uv \in E$ and $uw \in E$, with $v \neq w$ implies $c(uv) \neq c(uw)$
- $uv \in E$ implies $c(uv) \neq c(u)$




Total Coloring

Definition (k -total coloring)

A k -total coloring of $G = (V, E)$ is $c : V \cup E \rightarrow \{1, \dots, k\}$ such that:

- $uv \in E$ implies $c(u) \neq c(v)$
- $uv \in E$ and $uw \in E$, with $v \neq w$ implies $c(uv) \neq c(uw)$
- $uv \in E$ implies $c(uv) \neq c(u)$



Definition (total chromatic number)

$\chi_T(G)$ is the minimum k such that G admits a k -total coloring.

Famous bounds

Let $\Delta(G)$ denotes the maximum degree of G .

Famous bounds

Let $\Delta(G)$ denotes the maximum degree of G .

- Brook's Theorem (1941): $\chi(G) \leq \Delta(G)$ if G is not a complete graph or an odd cycle.

Famous bounds

Let $\Delta(G)$ denotes the maximum degree of G .

- Brook's Theorem (1941): $\chi(G) \leq \Delta(G)$ if G is not a complete graph or an odd cycle.
- Vizing's Theorem (1964): $\Delta(G) \leq \chi'(G) \leq \Delta(G) + 1$

Famous bounds

Let $\Delta(G)$ denotes the maximum degree of G .

- Brook's Theorem (1941): $\chi(G) \leq \Delta(G)$ if G is not a complete graph or an odd cycle.
- Vizing's Theorem (1964): $\Delta(G) \leq \chi'(G) \leq \Delta(G) + 1$
- Total coloring conjecture (Behzad, Vizing, ~1964):
 $\Delta(G) + 1 \leq \chi_T(G) \leq \Delta(G) + 2$

Algorithmic problems

k -VERTEX/EDGE/TOTAL COLORING:

- *Input*: a graph $G = (V, E)$
- *Output*: Does G admit a k -vertex/edge/total coloring?

Algorithmic problems

k -VERTEX/EDGE/TOTAL COLORING:

- *Input*: a graph $G = (V, E)$
 - *Output*: Does G admit a k -vertex/edge/total coloring?
- k -VERTEX COLORING is NP-complete for every fixed $k \geq 3$ (M.R. Garey and D.S. Johnson, 1979).

Algorithmic problems

k -VERTEX/EDGE/TOTAL COLORING:

- *Input*: a graph $G = (V, E)$
- *Output*: Does G admit a k -vertex/edge/total coloring?
- k -VERTEX COLORING is NP-complete for every fixed $k \geq 3$ (M.R. Garey and D.S. Johnson, 1979).
- k -EDGE COLORING is NP-complete for every fixed $k \geq 3$ (I. Holyer, 1981).

Algorithmic problems

k -VERTEX/EDGE/TOTAL COLORING:

- *Input*: a graph $G = (V, E)$
 - *Output*: Does G admit a k -vertex/edge/total coloring?
- k -VERTEX COLORING is NP-complete for every fixed $k \geq 3$ (M.R. Garey and D.S. Johnson, 1979).
 - k -EDGE COLORING is NP-complete for every fixed $k \geq 3$ (I. Holyer, 1981).
 - k -TOTAL COLORING is NP-complete for every fixed $k \geq 4$ (A. Sánchez-Arroyo, 1989).

Algorithmic problems

k -VERTEX/EDGE/TOTAL COLORING:

- *Input*: a graph $G = (V, E)$
- *Output*: Does G admit a k -vertex/edge/total coloring?
- k -VERTEX COLORING is NP-complete for every fixed $k \geq 3$ (M.R. Garey and D.S. Johnson, 1979).
- k -EDGE COLORING is NP-complete for every fixed $k \geq 3$ (I. Holyer, 1981).
- k -TOTAL COLORING is NP-complete for every fixed $k \geq 4$ (A. Sánchez-Arroyo, 1989).

ENUM- k -VERTEX/EDGE/TOTAL COLORING:

- *Input*: a graph $G = (V, E)$
- *Output*: Enumerate all the k -vertex/edge/total coloring of G .

Algorithmic problems

k -VERTEX/EDGE/TOTAL COLORING:

- *Input*: a graph $G = (V, E)$
- *Output*: Does G admit a k -vertex/edge/total coloring?
- k -VERTEX COLORING is NP-complete for every fixed $k \geq 3$ (M.R. Garey and D.S. Johnson, 1979).
- k -EDGE COLORING is NP-complete for every fixed $k \geq 3$ (I. Holyer, 1981).
- k -TOTAL COLORING is NP-complete for every fixed $k \geq 4$ (A. Sánchez-Arroyo, 1989).

ENUM- k -VERTEX/EDGE/TOTAL COLORING:

- *Input*: a graph $G = (V, E)$
- *Output*: Enumerate all the k -vertex/edge/total coloring of G .

Algorithmic point of view

What to do with an NP-complete problem?

Algorithmic point of view

What to do with an NP-complete problem?

- *approximation algorithm*: find in poly-time a result which is not worst than c time the optimal.
- *randomized algorithm*: find a result which is good w.h.p. or/and with running time correct w.h.p.
- ...

Algorithmic point of view

What to do with an NP-complete problem?

- *approximation algorithm*: find in poly-time a result which is not worst than c time the optimal.
- *randomized algorithm*: find a result which is good w.h.p. or/and with running time correct w.h.p.
- ...
- *parameterized algorithm*: find an algorithm with running time $f(k).poly(n)$ for some parameter k .

Algorithmic point of view

What to do with an NP-complete problem?

- *approximation algorithm*: find in poly-time a result which is not worst than c time the optimal.
- *randomized algorithm*: find a result which is good w.h.p. or/and with running time correct w.h.p.
- ...
- *parameterized algorithm*: find an algorithm with running time $f(k).poly(n)$ for some parameter k .
- **exact (exponential) algorithms**: find an algorithm with running time $O^*(c^n)$ ($= O(P(n)c^n)$)

Examples of exact algorithms

ENUM-3-EDGE COLORING:

- *Input*: a graph $G = (V, E)$ with $\Delta(G) \leq 3$.
- *Output*: Enumerate all the 3-vertex/edge/total coloring of G .

Examples of exact algorithms

ENUM-3-EDGE COLORING:

- *Input*: a graph $G = (V, E)$ with $\Delta(G) \leq 3$.
- *Output*: Enumerate all the 3-vertex/edge/total coloring of G .

The exhaustive algorithm:

- Try every possible colorings of the edge set and decide which colorings are proper.

Examples of exact algorithms

ENUM-3-EDGE COLORING:

- *Input*: a graph $G = (V, E)$ with $\Delta(G) \leq 3$.
- *Output*: Enumerate all the 3-vertex/edge/total coloring of G .

The exhaustive algorithm:

- Try every possible colorings of the edge set and decide which colorings are proper.

Time analysis:

$$O^*(3^m) = O^*(3^{\frac{3}{2}n}) = O^*(5.1962^n)$$

Examples of exact algorithms

ENUM-3-EDGE COLORING:

- *Input*: a graph $G = (V, E)$ with $\Delta(G) \leq 3$.
- *Output*: Enumerate all the 3-vertex/edge/total coloring of G .

The exhaustive algorithm:

- Try every possible colorings of the edge set and decide which colorings are proper.


Time analysis:

$$O^*(3^m) = O^*(3^{\frac{3}{2}n}) = O^*(5.1962^n)$$

Usual methods to improve: **branching algorithms** and dynamic programming.


Examples of exact algorithms

ENUM-3-EDGE COLORING: a 'naive' branching algorithms




Examples of exact algorithms

ENUM-3-EDGE COLORING: a 'naive' branching algorithms



Examples of exact algorithms

ENUM-3-EDGE COLORING: a 'naive' branching algorithms




Time analysis: $O^*(2^m) = O^*(2^{\frac{3}{2}n}) = O^*(2.8284^n)$

Polynomial space.

Examples of exact algorithms

ENUM-3-EDGE COLORING: a 'naive' branching algorithms



Time analysis: $O^*(2^m) = O^*(2^{\frac{3}{2}n}) = O^*(2.8284^n)$ min. nb of leaves

Polynomial space. min. height

Previous work: exact algorithms

What is the gain on the running time?

Previous work: exact algorithms

What is the gain on the running time?

In the same environment:

$$t = O^*(a^{n_1}) = O^*(b^{n_2}) \text{ with } a < b$$

We obtain: $n_1 \sim \frac{\log b}{\log a} n_2$

A multiplicative factor on the instance size of $\frac{\log b}{\log a} > 1$

Previous work: exact algorithms

What is the gain on the running time?

In the same environment:

$$t = O^*(a^{n_1}) = O^*(b^{n_2}) \text{ with } a < b$$

We obtain: $n_1 \sim \frac{\log b}{\log a} n_2$

A multiplicative factor on the instance size of $\frac{\log b}{\log a} > 1$

$$\text{Here: } \frac{\log 3^{\frac{3}{2}}}{\log 2^{\frac{3}{2}}} = 1.58$$

Previous work: exact algorithms

Main results in exact algorithms to color a graph:

- 3-VERTEX COLORING in $O^*(1.3289^n)$ (R. Beigel and D. Eppstein, 2005)

Previous work: exact algorithms

Main results in exact algorithms to color a graph:

- 3-VERTEX COLORING in $O^*(1.3289^n)$ (R. Beigel and D. Eppstein, 2005)
- MINIMUM VERTEX COLORING in $O^*(2^n)$ (A. Bjöklund, T. Husfeldt and M. Koivisto, 2006)

Previous work: exact algorithms

Main results in exact algorithms to color a graph:

- 3-VERTEX COLORING in $O^*(1.3289^n)$ (R. Beigel and D. Eppstein, 2005)
- MINIMUM VERTEX COLORING in $O^*(2^n)$ (A. Bjöklund, T. Husfeld and M. Koivisto, 2006)
- ENUM 3-VERTEX COLORING in $O^*(1.6259^n)$ (F. Fomin, S. Gasper and S. Saurabh, 2007)

Previous work: exact algorithms

Main results in exact algorithms to color a graph:

- 3-VERTEX COLORING in $O^*(1.3289^n)$ (R. Beigel and D. Eppstein, 2005)
- MINIMUM VERTEX COLORING in $O^*(2^n)$ (A. Bjöklund, T. Husfeld and M. Koivisto, 2006)
- ENUM 3-VERTEX COLORING in $O^*(1.6259^n)$ (F. Fomin, S. Gasper and S. Saurabh, 2007)
- 3-EDGE COLORING in $O^*(1.344^n)$ (L. Kowalik, 2009)

Previous work: exact algorithms

Main results in exact algorithms to color a graph:

- 3-VERTEX COLORING in $O^*(1.3289^n)$ (R. Beigel and D. Eppstein, 2005)
- MINIMUM VERTEX COLORING in $O^*(2^n)$ (A. Bjöklund, T. Husfeld and M. Koivisto, 2006)
- ENUM 3-VERTEX COLORING in $O^*(1.6259^n)$ (F. Fomin, S. Gasper and S. Saurabh, 2007)
- 3-EDGE COLORING in $O^*(1.344^n)$ (L. Kowalik, 2009)
- ENUM-3-EDGE COLORING in $O^*(1.5423^n)$ (P.A. Golovatch, D. Kratsch and J.F. Couturier, 2010)

Previous work: exact algorithms

Main results in exact algorithms to color a graph:

- 3-VERTEX COLORING in $O^*(1.3289^n)$ (R. Beigel and D. Eppstein, 2005)
- MINIMUM VERTEX COLORING in $O^*(2^n)$ (A. Bjöklund, T. Husfeld and M. Koivisto, 2006)
- ENUM 3-VERTEX COLORING in $O^*(1.6259^n)$ (F. Fomin, S. Gasper and S. Saurabh, 2007)
- 3-EDGE COLORING in $O^*(1.344^n)$ (L. Kowalik, 2009)
- ENUM-3-EDGE COLORING in $O^*(1.5423^n)$ (P.A. Golovatch, D. Kratsch and J.F. Couturier, 2010)
- ENUM-4-TOTAL COLORING in $O^*(3.0845^n)$ (P.A. Golovatch, D. Kratsch and J.F. Couturier, 2010)

Previous work: exact algorithms

Main results in exact algorithms to color a graph:

- 3-VERTEX COLORING in $O^*(1.3289^n)$ (R. Beigel and D. Eppstein, 2005)
- MINIMUM VERTEX COLORING in $O^*(2^n)$ (A. Bjöklund, T. Husfeld and M. Koivisto, 2006)
- ENUM 3-VERTEX COLORING in $O^*(1.6259^n)$ (F. Fomin, S. Gasper and S. Saurabh, 2007)
- 3-EDGE COLORING in $O^*(1.344^n)$ (L. Kowalik, 2009)
- ENUM-3-EDGE COLORING in $O^*(1.5423^n)$ (P.A. Golovatch, D. Kratsch and J.F. Couturier, 2010)
- ENUM-4-TOTAL COLORING in $O^*(3.0845^n)$ (P.A. Golovatch, D. Kratsch and J.F. Couturier, 2010)

Our results

Motivations:

- a useful technique for EDGE COLORING

Our results

Motivations:

- a useful technique for EDGE COLORING
- 3-EDGE COLORING more interesting

Our results

Motivations:

- a useful technique for EDGE COLORING
- 3-EDGE COLORING more interesting
- Gap: $\Omega(1.2822^n) \leq \text{ENUM-3-EDGE COLORING} \leq O^*(1.5423^n)$

Our results

Motivations:

- a useful technique for EDGE COLORING
- 3-EDGE COLORING more interesting
- Gap: $\Omega(1.2822^n) \leq \text{ENUM-3-EDGE COLORING} \leq O^*(1.5423^n)$
- $? \leq \text{ENUM-4-TOTAL COLORING} \leq O^*(3.0845^n)$

Our results

Motivations:

- a useful technique for EDGE COLORING
- 3-EDGE COLORING more interesting
- Gap: $\Omega(1.2822^n) \leq \text{ENUM-3-EDGE COLORING} \leq O^*(1.5423^n)$
- $? \leq \text{ENUM-4-TOTAL COLORING} \leq O^*(3.0845^n)$

Our results:

- ENUM-3-EDGE COLORING in $O^*(1.4142^n)$ and a sharp example (multi-graph). (factor on data size: 1.25)

Our results

Motivations:

- a useful technique for EDGE COLORING
- 3-EDGE COLORING more interesting
- Gap: $\Omega(1.2822^n) \leq \text{ENUM-3-EDGE COLORING} \leq O^*(1.5423^n)$
- $? \leq \text{ENUM-4-TOTAL COLORING} \leq O^*(3.0845^n)$

Our results:

- ENUM-3-EDGE COLORING in $O^*(1.4142^n)$ and a sharp example (multi-graph). (factor on data size: 1.25)
- ENUM-4-TOTAL COLORING in $O^*(2.8285^n)$. Example in $\Omega(2.3784^n)$.

Our results

Motivations:

- a useful technique for EDGE COLORING
- 3-EDGE COLORING more interesting
- Gap: $\Omega(1.2822^n) \leq \text{ENUM-3-EDGE COLORING} \leq O^*(1.5423^n)$
- $? \leq \text{ENUM-4-TOTAL COLORING} \leq O^*(3.0845^n)$

Our results:

- ENUM-3-EDGE COLORING in $O^*(1.4142^n)$ and a sharp example (multi-graph). (factor on data size: 1.25)
- ENUM-4-TOTAL COLORING in $O^*(2.8285^n)$. Example in $\Omega(2.3784^n)$.
- ENUM- k -EDGE COLORING in $O^*((k-1!)^{\frac{n}{2}})$ and a sharp example (multi-graph).

Our results

Motivations:

- a useful technique for EDGE COLORING
- 3-EDGE COLORING more interesting
- Gap: $\Omega(1.2822^n) \leq \text{ENUM-3-EDGE COLORING} \leq O^*(1.5423^n)$
- $? \leq \text{ENUM-4-TOTAL COLORING} \leq O^*(3.0845^n)$

Our results:

- ENUM-3-EDGE COLORING in $O^*(1.4142^n)$ and a sharp example (multi-graph). (factor on data size: 1.25)
- ENUM-4-TOTAL COLORING in $O^*(2.8285^n)$. Example in $\Omega(2.3784^n)$.
- ENUM- k -EDGE COLORING in $O^*((k-1!)^{\frac{n}{2}})$ and a sharp example (multi-graph).
- More precise bound on the number of 3-edge coloring of simple graph

Enumerating the 3-edge colorings

- 1 Introduction
 - Colorings
 - Algorithmic problems
 - Our results
- 2 Enumerating the 3-edge colorings of a cubic graph
 - The 3-edge colorings of a 3-regular graph
 - Turning the proof into algorithm
- 3 Extensions: k -edge colorings and the total colorings
 - k -edge colorings
 - A more precise bounds for the 3-edge coloring
 - Total coloring
- 4 Conclusion

ENUM-3-EDGE COLORING

We denote by $c_k(G)$ the number of k -edge-colorings of a graph G .


Here, we want to compute $c_3(G)$, G being a sub-cubic (multiple) graph.

ENUM-3-EDGE COLORING

We denote by $c_k(G)$ the number of k -edge-colorings of a graph G .

Here, we want to compute $c_3(G)$, G being a sub-cubic (multiple) graph.

We can assume that G is connected.




$$c_3(G) = c_3(G_1) \cdot c_3(G_2)$$

ENUM-3-EDGE COLORING

We denote by $c_k(G)$ the number of k -edge-colorings of a graph G .


Here, we want to compute $c_3(G)$, G being a sub-cubic (multiple) graph.

We can assume that G is connected.



$$c_3(G) = c_3(G_1) \cdot c_3(G_2)$$

We can assume that G is 2-(vertex) connected.



$$c_3(G) = \frac{1}{3} \cdot c_3(G_1) \cdot c_3(G_2)$$

Number of 3-edge colorings of 3-regular graphs

Lemma

Let C_n be the cycle of length n .

$$c_3(C_n) = \begin{cases} 2^n + 2, & \text{if } n \text{ is even,} \\ 2^n - 2, & \text{if } n \text{ is odd.} \end{cases}$$



Number of 3-edge colorings of 3-regular graphs

Lemma

Let C_n be the cycle of length n .

$$c_3(C_n) = \begin{cases} 2^n + 2, & \text{if } n \text{ is even,} \\ 2^n - 2, & \text{if } n \text{ is odd.} \end{cases}$$

Proof: By induction on n . True for $n = 2$ and $n = 3$.



$$c_3(C_n) = 2 \cdot c_3(C_{n-2}) +$$



Number of 3-edge colorings of 3-regular graphs

Lemma

Let C_n be the cycle of length n .

$$c_3(C_n) = \begin{cases} 2^n + 2, & \text{if } n \text{ is even,} \\ 2^n - 2, & \text{if } n \text{ is odd.} \end{cases}$$

Proof: By induction on n . True for $n = 2$ and $n = 3$.



$$c_3(C_n) = 2 \cdot c_3(C_{n-2}) + c_3(C_{n-1})$$



Number of 3-edge colorings of 3-regular graphs

We denote by n_i the number of vertices of degree i in G .

Theorem

Let G be a 2-connected subcubic graph. Then $c_3(G) \leq 3 \cdot 2^{n - \frac{n_3}{2}}$.

Number of 3-edge colorings of 3-regular graphs

We denote by n_i the number of vertices of degree i in G .


Theorem

Let G be a 2-connected subcubic graph. Then $c_3(G) \leq 3 \cdot 2^{n - \frac{n_3}{2}}$.

Proof: If G is a cycle, it is true.

Otherwise, let v_1 and v_n be two vertices of degree 3 and consider v_1, \dots, v_n an *st-ordering* of G :

for all $1 < i < n$, $d(v_i)_{\{v_1, \dots, v_{i-1}\}} \geq 1$ and $d(v_i)_{\{v_{i+1}, \dots, v_n\}} \geq 1$.
(always exists: Lempel et al., 1967)




Number of 3-edge colorings of 3-regular graphs

Theorem

Let G be a 2-connected subcubic graph. Then $c_3(G) \leq 3 \cdot 2^{n - \frac{n_3}{2}}$.

Proof:



Orient from left to right.

$$A^+ = \{v : d^+(v) = 2 \text{ and } d^-(v) = 1\}$$

$$A^- = \{v : d^+(v) = 1 \text{ and } d^-(v) = 2\}$$


$$\text{We have } |A^+| = |A^-| = (n_3 - 2)/2$$

Number of 3-edge colorings of 3-regular graphs

Theorem

Let G be a 2-connected subcubic graph. Then $c_3(G) \leq 3 \cdot 2^{n - \frac{n_3}{2}}$.

Proof:



Orient from left to right.

$$A^+ = \{v : d^+(v) = 2 \text{ and } d^-(v) = 1\}$$

$$A^- = \{v : d^+(v) = 1 \text{ and } d^-(v) = 2\}$$

$$\text{We have } |A^+| = |A^-| = (n_3 - 2)/2$$

Number of 3-edge colorings of 3-regular graphs

Theorem

Let G be a 2-connected subcubic graph. Then $c_3(G) \leq 3 \cdot 2^{n - \frac{n_3}{2}}$.

Proof: We have $|A^+| = |A^-| = (n_3 - 2)/2$

Denote by c_i the number of partial 3-edge colorings of arcs with tail in $\{v_1, \dots, v_i\}$.

Number of 3-edge colorings of 3-regular graphs


Theorem


Let G be a 2-connected subcubic graph. Then $c_3(G) \leq 3 \cdot 2^{n - \frac{n_3}{2}}$.


Proof: We have $|A^+| = |A^-| = (n_3 - 2)/2$

Denote by c_i the number of partial 3-edge colorings of arcs with tail in $\{v_1, \dots, v_i\}$.

- $c_1 = 6$

- if $v_i \in A^-$  $c_i \leq c_{i-1}$

- if $v_i \in A^+$  $c_i \leq 2c_{i-1}$

- if $d^+(v_i) = d^-(v_i) = 1$  $c_i \leq 2c_{i-1}$

Number of 3-edge colorings of 3-regular graphs


Theorem


Let G be a 2-connected subcubic graph. Then $c_3(G) \leq 3 \cdot 2^{n - \frac{n_3}{2}}$.


Proof: We have $|A^+| = |A^-| = (n_3 - 2)/2$

Denote by c_i the number of partial 3-edge colorings of arcs with tail in $\{v_1, \dots, v_i\}$.

- $c_1 = 6$

- if $v_i \in A^-$  $c_i \leq c_{i-1}$

- if $v_i \in A^+$  $c_i \leq 2c_{i-1}$

- if $d^+(v_i) = d^-(v_i) = 1$  $c_i \leq 2c_{i-1}$

$$c_3(G) \leq 6 \cdot 2^{(n-2) - (n_3-2)/2} = 3 \cdot 2^{n - \frac{n_3}{2}}$$



Number of 3-edge colorings of 3-regular graphs


Theorem

Let G be a 2-connected subcubic graph. Then $c_3(G) \leq 3 \cdot 2^{n - \frac{n_3}{2}}$.

Corollary

Let G be a cubic graph, then $c_3(G) \leq 3 \cdot 2^{\frac{n}{2}}$.

This is sharp for multi-graphs:



ENUM-3-EDGE COLORING:

Corollary (Solving ENUM-3-EDGE COLORING:)

There exists a branching algorithm with running time $O^(2^{\frac{n}{2}}) = O^*(1.4143^n)$ and polynomial space to solve ENUM-3-EDGE COLORING.*

ENUM-3-EDGE COLORING:


Corollary (Solving ENUM-3-EDGE COLORING:)

There exists a branching algorithm with running time $O^(2^{\frac{n}{2}}) = O^*(1.4143^n)$ and polynomial space to solve ENUM-3-EDGE COLORING.*

Proof:

- If $\Delta \geq 4$ answer 0.
- If the graph is not connected enough, divide the instance.
- Otherwise, run a branching algorithm.

ENUM-3-EDGE COLORING:



Extensions

- 1 Introduction
 - Colorings
 - Algorithmic problems
 - Our results
- 2 Enumerating the 3-edge colorings of a cubic graph
 - The 3-edge colorings of a 3-regular graph
 - Turning the proof into algorithm
- 3 Extensions: k -edge colorings and the total colorings
 - k -edge colorings
 - A more precise bounds for the 3-edge coloring
 - Total coloring
- 4 Conclusion

Number of k -edge colorings of k -regular graphs

We denote by c_k the number of k -edge colorings of G .

Theorem

Let G be a k -regular graph. Then $c_k(G) \leq k \cdot (k-1!)^{\frac{n}{2}}$.

Number of k -edge colorings of k -regular graphs

We denote by c_k the number of k -edge colorings of G .

Theorem

Let G be a k -regular graph. Then $c_k(G) \leq k \cdot (k-1!)^{\frac{n}{2}}$.

Proof: G is 2-connected,


Number of k -edge colorings of k -regular graphs

We denote by c_k the number of k -edge colorings of G .

Theorem

Let G be a k -regular graph. Then $c_k(G) \leq k \cdot (k-1!)^{\frac{n}{2}}$.

Proof: G is 2-connected, consider v_1, \dots, v_n an st -ordering of G :




Number of k -edge colorings of k -regular graphs

We denote by c_k the number of k -edge colorings of G .

Theorem

Let G be a k -regular graph. Then $c_k(G) \leq k \cdot (k-1!)^{\frac{n}{2}}$.

Proof: G is 2-connected, consider v_1, \dots, v_n an st -ordering of G :




$A_i = \{v : d^+(v) = i\}$ and $a_i = |A_i|$ for $i = 1, \dots, k-1$

Then, $c_k(G) \leq \prod_{i=1}^{n-1} d^+(v_i)! \leq k! \prod_{i=1}^{k-1} (i!)^{a_i}$
 under $\sum_{i=1}^{k-1} a_i = n-2$ and $\sum_{i=1}^{k-1} i \cdot a_i = k(n-2)/2$

□

Number of k -edge colorings of k -regular graphs

Also sharp for multi-graphs:



ENUM- k -EDGE COLORING:

Corollary (Solving ENUM-3-EDGE COLORING:)


There exists a branching algorithm with running time $O^((k-1!)^{\frac{n}{2}})$ and polynomial space to solve ENUM- k -EDGE COLORING.*

ENUM- k -EDGE COLORING:

Corollary (Solving ENUM-3-EDGE COLORING:)

There exists a branching algorithm with running time $O^((k-1!)^{\frac{n}{2}})$ and polynomial space to solve ENUM- k -EDGE COLORING.*

Proof:




At most $\prod_{i=1}^{n-1} d^+(v_i)! = O^*((k-1!)^{\frac{n}{2}})$ leaves

Number of 3-edge colorings for simple graphs

Corollary

Let G be a cubic graph, then $c_3(G) \leq 3 \cdot 2^{\frac{n}{2}}$.

The sharp example for the number of 3-edge coloring of cubic graphs:




Number of 3-edge colorings for simple graphs

Can we improve a lot?

Number of 3-edge colorings for simple graphs

Can we improve a lot? Not really: let H_n be:




Lemma (c_3 of the ladder graph)

$$c_3(H_n) = \begin{cases} 2^{n/2} + 8, & \text{if } n/2 \text{ is even,} \\ 2^{n/2} - 2, & \text{if } n/2 \text{ is odd.} \end{cases}$$

Number of 3-edge colorings for simple graphs

Can we improve a lot? Not really: let H_n be:




Lemma (c_3 of the ladder graph)

$$c_3(H_n) = \begin{cases} 2^{n/2} + 8, & \text{if } n/2 \text{ is even,} \\ 2^{n/2} - 2, & \text{if } n/2 \text{ is odd.} \end{cases}$$

Number of 3-edge colorings for simple graphs

Can we improve a lot? Not really: let H_n be:




Lemma (c_3 of the ladder graph)

$$c_3(H_n) = \begin{cases} 2^{n/2} + 8, & \text{if } n/2 \text{ is even,} \\ 2^{n/2} - 2, & \text{if } n/2 \text{ is odd.} \end{cases}$$

Number of 3-edge colorings for simple graphs

Can we improve a lot? Not really: let H_n be:




Lemma (c_3 of the ladder graph)

$$c_3(H_n) = \begin{cases} 2^{n/2} + 8, & \text{if } n/2 \text{ is even,} \\ 2^{n/2} - 2, & \text{if } n/2 \text{ is odd.} \end{cases}$$

Number of 3-edge colorings for simple graphs

Can we improve a lot? Not really: let H_n be:




Lemma (c_3 of the ladder graph)

$$c_3(H_n) = \begin{cases} 2^{n/2} + 8, & \text{if } n/2 \text{ is even,} \\ 2^{n/2} - 2, & \text{if } n/2 \text{ is odd.} \end{cases}$$

Number of 3-edge colorings for simple graphs

Can we improve a lot? Not really: let H_n be:




Lemma (c_3 of the ladder graph)

$$c_3(H_n) = \begin{cases} 2^{n/2} + 8, & \text{if } n/2 \text{ is even,} \\ 2^{n/2} - 2, & \text{if } n/2 \text{ is odd.} \end{cases}$$

Number of 3-edge colorings for simple graphs

Can we improve a lot? Not really: let H_n be:




Lemma (c_3 of the ladder graph)

$$c_3(H_n) = \begin{cases} 2^{n/2} + 8, & \text{if } n/2 \text{ is even,} \\ 2^{n/2} - 2, & \text{if } n/2 \text{ is odd.} \end{cases}$$

Number of 3-edge colorings for simple graphs

Can we improve a lot? Not really: let M_n be:



Lemma (c_3 of the Möbius ladder graph)

$$c_3(M_n) = \begin{cases} 2^{n/2} + 2, & \text{if } n/2 \text{ is even,} \\ 2^{n/2} + 4, & \text{if } n/2 \text{ is odd.} \end{cases}$$

Number of 3-edge colorings for simple graphs

So, for simple cubic graphs, we have:

$$2^{\frac{n}{2}} < \max\{c_3(G) : G \text{ cubic graph on } n \text{ vertices}\} \leq 3.2^{\frac{n}{2}}$$

Number of 3-edge colorings for simple graphs

So, for simple cubic graphs, we have:

$$2^{\frac{n}{2}} < \max\{c_3(G) : G \text{ cubic graph on } n \text{ vertices}\} \leq \cancel{3 \cdot 2^{\frac{n}{2}}} \frac{9}{4} \cdot 2^{\frac{n}{2}}$$

Theorem

Let G be a connected simple cubic graph. Then $c_3(G) \leq \frac{9}{4} \cdot 2^{n-\frac{n_3}{2}}$.

Number of 3-edge colorings for simple graphs


So, for simple cubic graphs, we have:

$$2^{\frac{n}{2}} < \max\{c_3(G) : G \text{ cubic graph on } n \text{ vertices}\} \leq \cancel{3 \cdot 2^{\frac{n}{2}}} \frac{9}{4} \cdot 2^{\frac{n}{2}}$$

Theorem

Let G be a connected simple cubic graph. Then $c_3(G) \leq \frac{9}{4} \cdot 2^{n-\frac{n_3}{2}}$.

Proof:



We have $\frac{1}{3} \cdot c_3(C_{j-i+1})/2^{j-i} \cdot 3 \cdot 2^{\frac{n}{2}}$ colorings.

4-total colorings

Let $c_4^T(G)$ be the number of 4-total colorings of G .

4-total colorings

Let $c_4^T(G)$ be the number of 4-total colorings of G .

Theorem

Let G be a (2-)connected cubic graph. Then $c_4^T(G) \leq 3 \cdot 2^{\frac{3n}{2}}$.


4-total colorings

Let $c_4^T(G)$ be the number of 4-total colorings of G .

Theorem

Let G be a (2-)connected cubic graph. Then $c_4^T(G) \leq 3 \cdot 2^{\frac{3n}{2}}$.

Proof:



Orient from left to right.

$$A^+ = \{v : d^+(v) = 2 \text{ and } d^-(v) = 1\}$$

$$A^- = \{v : d^+(v) = 1 \text{ and } d^-(v) = 2\}$$

$$\text{We have } |A^+| = |A^-| = (n-2)/2$$

4-total colorings

Theorem

Let G be a $(2-)$ connected cubic graph. Then $c_4^T(G) \leq 3 \cdot 2^{\frac{3n}{2}}$.

Proof: We have $|A^+| = |A^-| = (n-2)/2$

Denote by c_i the number of partial 4-total colorings of vertices and arcs with tail in $\{v_1, \dots, v_i\}$.

4-total colorings

Theorem

Let G be a $(2-)$ connected cubic graph. Then $c_4^T(G) \leq 3 \cdot 2^{\frac{3n}{2}}$.

Proof: We have $|A^+| = |A^-| = (n-2)/2$

Denote by c_i the number of partial 4-total colorings of vertices and arcs with tail in $\{v_1, \dots, v_i\}$.

- $c_1 = 24$

4-total colorings


Theorem

Let G be a $(2-)$ connected cubic graph. Then $c_4^T(G) \leq 3 \cdot 2^{\frac{3n}{2}}$.

Proof: We have $|A^+| = |A^-| = (n-2)/2$

Denote by c_i the number of partial 4-total colorings of vertices and arcs with tail in $\{v_1, \dots, v_i\}$.

- $c_1 = 24$

- if $v_i \in A^-$  $c_i \leq 2c_{i-1}$

4-total colorings


Theorem


Let G be a $(2-)$ connected cubic graph. Then $c_4^T(G) \leq 3 \cdot 2^{\frac{3n}{2}}$.

Proof: We have $|A^+| = |A^-| = (n-2)/2$

Denote by c_i the number of partial 4-total colorings of vertices and arcs with tail in $\{v_1, \dots, v_i\}$.

- $c_1 = 24$

- if $v_i \in A^-$  $c_i \leq 2c_{i-1}$

- if $v_i \in A^+$  $c_i \leq 4c_{i-1}$

4-total colorings

Theorem


Let G be a $(2-)$ connected cubic graph. Then $c_4^T(G) \leq 3 \cdot 2^{\frac{3n}{2}}$.

Proof: We have $|A^+| = |A^-| = (n-2)/2$

Denote by c_i the number of partial 4-total colorings of vertices and arcs with tail in $\{v_1, \dots, v_i\}$.

- $c_1 = 24$

- if $v_i \in A^-$  $c_i \leq 2c_{i-1}$

- if $v_i \in A^+$  $c_i \leq 4c_{i-1}$

Hence:

$$c_4^T(G) \leq 24 \cdot 2^{\frac{(n-2)}{2}} \cdot 4^{\frac{(n-2)}{2}} = 3 \cdot 2^{\frac{3n}{2}}$$

□

ENUM-4-TOTAL COLORING:

Corollary (Solving ENUM-3-EDGE COLORING:)

There exists a branching algorithm with running time $O^(2^{\frac{3n}{2}}) = O^*(2.8285^n)$ and polynomial space to solve ENUM-4-TOTAL COLORING.*

ENUM-4-TOTAL COLORING:

Corollary (Solving ENUM-3-EDGE COLORING:)

There exists a branching algorithm with running time $O^(2^{\frac{3n}{2}}) = O^*(2.8285^n)$ and polynomial space to solve ENUM-4-TOTAL COLORING.*

ENUM-4-TOTAL COLORING:

A lower bound on c_4^T , the nb of 4-total colorings:

ENUM-4-TOTAL COLORING:

A lower bound on c_4^T , the nb of 4-total colorings:

Lemma

Let T be a binary tree (degree = 1 or 3) on n vertices, $c_4^T(T) = 3 \cdot 2^{\frac{3n}{2}}$.

ENUM-4-TOTAL COLORING:

A lower bound on c_4^T , the nb of 4-total colorings:

Lemma

Let T be a binary tree (degree = 1 or 3) on n vertices, $c_4^T(T) = 3 \cdot 2^{\frac{3n}{2}}$.

Proof: by induction:

ENUM-4-TOTAL COLORING:

A lower bound on c_4^T , the nb of 4-total colorings:

Lemma

Let T be a binary tree (degree = 1 or 3) on n vertices, $c_4^T(T) = 3 \cdot 2^{\frac{3n}{2}}$.

Proof: by induction:

- $c_4^T(K_{1,3}) = 4 \cdot 3! \cdot 2^3 = 3 \cdot 2^6 = 3 \cdot 2^{\frac{3 \cdot 4}{2}}$

ENUM-4-TOTAL COLORING:


A lower bound on c_4^T , the nb of 4-total colorings:

Lemma

Let T be a binary tree (degree = 1 or 3) on n vertices, $c_4^T(T) = 3 \cdot 2^{\frac{3n}{2}}$.

Proof: by induction:

- $c_4^T(K_{1,3}) = 4 \cdot 3! \cdot 2^3 = 3 \cdot 2^6 = 3 \cdot 2^{\frac{3 \cdot 4}{2}}$
- $c_4^T(T) = 8 \cdot c_4^T(T \setminus \{x, y\})$



ENUM-4-TOTAL COLORING:


A lower bound on c_4^T , the nb of 4-total colorings:

Lemma

Let T be a binary tree (degree = 1 or 3) on n vertices, $c_4^T(T) = 3 \cdot 2^{\frac{3n}{2}}$.

Proof: by induction:

- $c_4^T(K_{1,3}) = 4 \cdot 3! \cdot 2^3 = 3 \cdot 2^6 = 3 \cdot 2^{\frac{3 \cdot 4}{2}}$
- $c_4^T(T) = 8 \cdot c_4^T(T \setminus \{x, y\})$




ENUM-4-TOTAL COLORING:

Lemma

Let T_n be a binary tree on n vertices + parallel edges between twin leaves,
 $c_4^T(T_n) = \frac{3}{\sqrt{2}} \cdot 2^{\frac{5n}{4}}$.

Proof:




$$n = p - 2 + p + 2p$$

ENUM-4-TOTAL COLORING:

Lemma

Let T_n be a binary tree on n vertices + parallel edges between twin leaves,
 $c_4^T(T_n) = \frac{3}{\sqrt{2}} \cdot 2^{\frac{5n}{4}}$.

Proof:




$$c_4^T(T') = 3 \cdot 2^{\frac{3(2p-2)}{2}}$$

ENUM-4-TOTAL COLORING:

Lemma

Let T_n be a binary tree on n vertices + parallel edges between twin leaves,
 $c_4^T(T_n) = \frac{3}{\sqrt{2}} \cdot 2^{\frac{5n}{4}}$.

Proof:




$$c_4^T(T') = 3 \cdot 2^{\frac{3(2p-2)}{2}}$$

ENUM-4-TOTAL COLORING:

Lemma

Let T_n be a binary tree on n vertices + parallel edges between twin leaves,
 $c_4^T(T_n) = \frac{3}{\sqrt{2}} \cdot 2^{\frac{5n}{4}}$.

Proof:




$$c_4^T(T') = 3 \cdot 2^{\frac{3(2p-2)}{2}} \cdot 4^p$$

ENUM-4-TOTAL COLORING:

Lemma

Let T_n be a binary tree on n vertices + parallel edges between twin leaves,
 $c_4^T(T_n) = \frac{3}{\sqrt{2}} \cdot 2^{\frac{5n}{4}}$.

Proof:



$$c_4^T(T') = 3 \cdot 2^{\frac{3(2p-2)}{2}} \cdot 4^p = 3 \cdot 2^{5p-3} = 3 \cdot 2^{\frac{5n}{4} - \frac{1}{2}}$$

Conclusion

- 1 Introduction
 - Colorings
 - Algorithmic problems
 - Our results
- 2 Enumerating the 3-edge colorings of a cubic graph
 - The 3-edge colorings of a 3-regular graph
 - Turning the proof into algorithm
- 3 Extensions: k -edge colorings and the total colorings
 - k -edge colorings
 - A more precise bounds for the 3-edge coloring
 - Total coloring
- 4 Conclusion

Some (hard?) open questions

- MINIMUM VERTEX COLORING in $O^*(2^n)$ (A. Bjöklund, T. Husfeld and M. Koivisto, 2006)

Some (hard?) open questions

- MINIMUM VERTEX COLORING in $O^*(2^n)$ (A. Bjöklund, T. Husfeld and M. Koivisto, 2006)

Is there a $O^*((2 - \epsilon)^n)$ algorithm for some $\epsilon > 0$

Some (hard?) open questions

- MINIMUM VERTEX COLORING in $O^*(2^n)$ (A. Bjöklund, T. Husfeldt and M. Koivisto, 2006)
Is there a $O^*((2 - \epsilon)^n)$ algorithm for some $\epsilon > 0$
- MINIMUM EDGE COLORING in $O^*(2^m)$

Some (hard?) open questions

- MINIMUM VERTEX COLORING in $O^*(2^n)$ (A. Bjöklund, T. Husfeldt and M. Koivisto, 2006)
Is there a $O^*((2 - \epsilon)^n)$ algorithm for some $\epsilon > 0$
- MINIMUM EDGE COLORING in $O^*(2^m)$
Is there a $O^*(2^n)$ algorithm

Some (hard?) open questions

- MINIMUM VERTEX COLORING in $O^*(2^n)$ (A. Bjöklund, T. Husfeldt and M. Koivisto, 2006)
Is there a $O^*((2 - \epsilon)^n)$ algorithm for some $\epsilon > 0$
- MINIMUM EDGE COLORING in $O^*(2^m)$
Is there a $O^*(2^n)$ algorithm
- 3-VERTEX COLORING in $O^*(1.3289^n)$ (R. Beigel and D. Eppstein, 2005)

Some (hard?) open questions

- MINIMUM VERTEX COLORING in $O^*(2^n)$ (A. Bjöklund, T. Husfeldt and M. Koivisto, 2006)
Is there a $O^*((2 - \epsilon)^n)$ algorithm for some $\epsilon > 0$
- MINIMUM EDGE COLORING in $O^*(2^m)$
Is there a $O^*(2^n)$ algorithm
- 3-VERTEX COLORING in $O^*(1.3289^n)$ (R. Beigel and D. Eppstein, 2005)
Is it possible to do better?

Some (hard?) open questions

- MINIMUM VERTEX COLORING in $O^*(2^n)$ (A. Bjöklund, T. Husfeldt and M. Koivisto, 2006)
Is there a $O^*((2 - \epsilon)^n)$ algorithm for some $\epsilon > 0$
- MINIMUM EDGE COLORING in $O^*(2^m)$
Is there a $O^*(2^n)$ algorithm
- 3-VERTEX COLORING in $O^*(1.3289^n)$ (R. Beigel and D. Eppstein, 2005)
Is it possible to do better?
Under ETH, there is no $O^*(2^{o(n)})$ algorithm for 3-VERTEX COLORING (D. Lokshtanov, D. Marx, S. Saurabh, 2012), several scenarios...

Some (less hard?) open questions

- ENUM-3-EDGE COLORING in $O^*(1.4142^n)$ and a sharp example (multi-graph).

Some (less hard?) open questions

- ENUM-3-EDGE COLORING in $O^*(1.4142^n)$ and a sharp example (multi-graph).

Is it possible to fix

$$c_3(n) = \max\{c_3(G) : G \text{ simple graph on } n \text{ vertices}\}$$

We know that $2^{\frac{n}{2}} < c_3(n) \leq \frac{9}{4} \cdot 2^{\frac{n}{2}}$

Some (less hard?) open questions

- ENUM-3-EDGE COLORING in $O^*(1.4142^n)$ and a sharp example (multi-graph).

Is it possible to fix

$$c_3(n) = \max\{c_3(G) : G \text{ simple graph on } n \text{ vertices}\}$$

We know that $2^{\frac{n}{2}} < c_3(n) \leq \frac{9}{4} \cdot 2^{\frac{n}{2}}$

- ENUM-4-TOTAL COLORING in $O^*(2.8285^n)$. Example in $\Omega(2.3784^n)$.

Some (less hard?) open questions

- ENUM-3-EDGE COLORING in $O^*(1.4142^n)$ and a sharp example (multi-graph).

Is it possible to fix

$$c_3(n) = \max\{c_3(G) : G \text{ simple graph on } n \text{ vertices}\}$$

We know that $2^{\frac{n}{2}} < c_3(n) \leq \frac{9}{4} \cdot 2^{\frac{n}{2}}$

- ENUM-4-TOTAL COLORING in $O^*(2.8285^n)$. Example in $\Omega(2.3784^n)$.

Is it possible to find the algorithm with best running time for ENUM-4-TOTAL COLORING.

Thank you for your attention !